

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1623PAZ

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page for STN Seminar Schedule - N. America
NEWS 2 AUG 10 Time limit for inactive STN sessions doubles to 40
minutes
NEWS 3 AUG 18 COMPENDEX indexing changed for the Corporate Source
(CS) field
NEWS 4 AUG 24 ENCOMPLIT/ENCOMPLIT2 reloaded and enhanced
NEWS 5 AUG 24 CA/CAPLUS enhanced with legal status information for
U.S. patents
NEWS 6 SEP 09 50 Millionth Unique Chemical Substance Recorded in
CAS REGISTRY
NEWS 7 SEP 11 WPIDS, WPINDEX, and WPIX now include Japanese FTERM
thesaurus
NEWS 8 OCT 21 Derwent World Patents Index Coverage of Indian and
Taiwanese Content Expanded
NEWS 9 OCT 21 Derwent World Patents Index enhanced with human
translated claims for Chinese Applications and
Utility Models
NEWS 10 NOV 23 Addition of SCAN format to selected STN databases
NEWS 11 NOV 23 Annual Reload of IFI Databases
NEWS 12 DEC 01 FRFULL Content and Search Enhancements
NEWS 13 DEC 01 DGENE, USGENE, and PCTGEN: new percent identity
feature for sorting BLAST answer sets
NEWS 14 DEC 02 Derwent World Patent Index: Japanese FI-TERM
thesaurus added
NEWS 15 DEC 02 PCTGEN enhanced with patent family and legal status
display data from INPADOCDB
NEWS 16 DEC 02 USGENE: Enhanced coverage of bibliographic and
sequence information
NEWS 17 DEC 21 New Indicator Identifies Multiple Basic Patent
Records Containing Equivalent Chemical Indexing
in CA/CAPLUS
NEWS 18 JAN 12 Match STN Content and Features to Your Information
Needs, Quickly and Conveniently
NEWS 19 JAN 25 Annual Reload of MEDLINE database
NEWS 20 FEB 16 STN Express Maintenance Release, Version 8.4.2, Is
Now Available for Download
NEWS 21 FEB 16 Derwent World Patents Index (DWPI) Revises Indexing
of Author Abstracts
NEWS 22 FEB 16 New FASTA Display Formats Added to USGENE and PCTGEN
NEWS 23 FEB 16 INPADOCDB and INPAFAMDB Enriched with New Content
and Features
NEWS 24 FEB 16 INSPEC Adding Its Own IPC codes and Author's E-mail
Addresses

NEWS EXPRESS FEBRUARY 15 10 CURRENT WINDOWS VERSION IS V8.4.2,
AND CURRENT DISCOVER FILE IS DATED 15 JANUARY 2010.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN customer agreement. This agreement limits use to scientific research. Use for software development or design, implementation of commercial gateways, or use of CAS and STN data in the building of commercial products is prohibited and may result in loss of user privileges and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 08:11:06 ON 16 MAR 2010

=> logoff hold

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.22	0.22

SESSION WILL BE HELD FOR 120 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 08:11:23 ON 16 MAR 2010

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1623PAZ

PASSWORD:

***** RECONNECTED TO STN INTERNATIONAL *****
SESSION RESUMED IN FILE 'HOME' AT 08:46:24 ON 16 MAR 2010
FILE 'HOME' ENTERED AT 08:46:24 ON 16 MAR 2010

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.22	0.22

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	1.10	1.10

FILE 'REGISTRY' ENTERED AT 08:48:41 ON 16 MAR 2010
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2010 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 15 MAR 2010 HIGHEST RN 1210111-73-1
DICTIONARY FILE UPDATES: 15 MAR 2010 HIGHEST RN 1210111-73-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 8, 2010.



Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=>

Uploading C:\Documents and Settings\PZucker\My Documents\Examination Auxillary files\10588286\10588286 core intermediate.str



```
chain nodes :
1  2  3  4  5  6  7  8  9  10 12 13
chain bonds :
1-2  2-3  2-10 3-4  3-5  4-8  4-9  5-6  5-7  6-12  8-13
exact/norm bonds :
1-2  4-8  4-9  5-6  5-7  6-12  8-13
exact bonds :
2-3  2-10  3-4  3-5
```

```
Hydrogen count :
2:>= minimum 1  3:>= minimum 1  10:>= minimum 3
Match level :
1:CLASS  2:CLASS  3:CLASS  4:CLASS  5:CLASS  6:CLASS  7:CLASS  8:CLASS  9:CLASS
10:CLASS 12:CLASS 13:CLASS
Generic attributes :
1:
Saturation           : Saturated


Element Count :
Node 1: Limited
C,C2-6
```

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using SIN Express query preparation.

=> search l1 sss sam

SAMPLE SEARCH INITIATED 08:49:14 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 9296 TO ITERATE

21.5% PROCESSED 2000 ITERATIONS

3 ANSWERS

INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 180140 TO 191700

PROJECTED ANSWERS: 54 TO 502


L2 3 SEA SSS SAM L1

=> d scan

L2 3 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN

IN Propanedioic acid, 2-(1-methyl-2-oxopropyl)-, 1,3-dimethyl ester

MF C9 H14 O5



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):3

L2 3 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN

IN Propanedioic acid, 2-(3-[1,1'-biphenyl]-4-yl-1-methyl-3-oxopropyl)-,


1,3-diethyl ester

MF C23 H26 O5



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L2 3 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-(3-fluoro-1-methylpropyl)-, 1,3-diethyl ester
 MF C11 H19 F O4



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

ALL ANSWERS HAVE BEEN SCANNED

=> save temp rawmalon8s/a
 ENTER L#, L# RANGE, ALL, OR (END):12
 ANSWER SET L2 HAS BEEN SAVED AS 'RAWMALON8S/A'

=> search l1 sss full
 FULL SEARCH INITIATED 08:51:08 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 182998 TO ITERATE

100.0% PROCESSED 182998 ITERATIONS
 SEARCH TIME: 00.00.02


274 ANSWERS

L3 274 SEA SSS FUL L1

=> d scan

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-[(1R)-3-[(1,1-dimethylethyl)thio]-1-methyl-3-oxopropyl]-, 1,3-dimethyl ester
 MF C13 H22 O5 S

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN

IN Propanedioic acid, 2-[5-[2-(acetyloxy)-1-ethyl-5-oxocyclopentyl]-1-methyl-3-oxopentyl]-, 1,3-dimethyl ester

MF C20 H30 O8




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN

IN Propanedioic acid, 2-[(1R)-1-methylbutyl]-, 1,3-dimethyl ester

MF C10 H18 O4

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN

IN Iron, tricarbonyl[dimethyl [2-[(2,3,4,5-η)-4-methoxy-2,4-cyclohexadien-1-ylidene]-1-methylpropyl]propanedioate]-, stereoisomer (9CI)


MF C19 H22 Fe O8

CI CCS




L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, (1-methylpentyl)-, diethyl ester, (-)- (9CI)
 MF C13 H24 O4

Rotation (-).



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN INDEX NAME NOT YET ASSIGNED
 MF C19 H36 O4




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN

IN 1,1,4,4-Butanetetra-carboxylic acid, 2,3-dimethyl-, tetramethyl ester,
 (R*,R*)- (9CI)
 MF C14 H22 O8


Relative stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN 1,1,6-Hexanetricarboxylic acid, 6-(benzoyloxy)-2-methyl-, 1,1-dimethyl
 ester, [R-(R*,R*)]- (9CI)
 MF C19 H24 O8

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN INDEX NAME NOT YET ASSIGNED
 MF C17 H24 O4




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
IN Propanedioic acid, 2-[3-[[1,1-dimethylethyl]dimethylsilyloxy]-1-
methylbutyl]-, 1,3-diethyl ester
MF C18 H36 O5 Si



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
IN Propanedioic acid, 2-[1-methyl-3-(1-methylethoxy)propyl]-, 1,3-diethyl
ester
MF C14 H26 O5




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10


L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
IN Propanedioic acid, 2-[(1R,3E)-4,4,4-trifluoro-3-[(2S)-2-(methoxymethyl)-1-
pyrrolidinyl]imino]-1-methylbutyl]-, 1,3-dimethyl ester
MF C16 H25 F3 N2 O5

Absolute stereochemistry.
Double bond geometry as shown.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN 2,2,4-Pentanetricarboxylic acid, 4-cyano-3-methyl-, 1,2,4-triethyl ester
 MF C14 H21 N O6



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-(3-cyano-1-methylpropyl)-, 1,3-diethyl ester
 MF C12 H19 N O4



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-[(1R,2R)-2-(4-methoxyphenyl)-1-methylpropyl]-,
 1,3-diethyl ester, rel-
 MF C18 H26 O5


Relative stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, [(1R,2R)-1-methyl-2-phenylpropyl]-, diethyl ester, rel-
 (9CI)
 MF C17 H24 O4

Relative stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-[1-methyl-2-(phenylmethoxy)propyl]-, 1,3-diethyl
 ester
 MF C18 H26 O5



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-[(1R,3S)-3-(dimethylphenylsilyl)-1,4-dimethylpentyl]-
 , 1,3-dimethyl ester, rel-
 MF C20 H32 O4 Si

Relative stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-(1,2-dimethyl-3-oxopropyl)-, 1,3-diethyl ester
 MF C12 H20 O5




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-(1-methyl-3-oxopentyl)-, 1,3-dimethyl ester
 MF C11 H18 O5



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L3 274 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN
 IN Propanedioic acid, 2-(3-amino-2-cyano-1-methyl-3-oxopropyl)-, 1,3-diethyl ester
 MF C12 H18 N2 O5



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):end

```
=> save temp l3 rawmalon8s/a
'RAWMALON8S/A' IN USE
A single name cannot be used for two saved items at the same time.
Enter "Y" if you wish to replace the current saved name with a new
definition. Enter "N" if the current saved definition must be
preserved. You may then reenter the SAVE command with a different
saved name. Enter "DISPLAY SAVED" at an arrow prompt (=>) to see a
list of your currently defined saved names.
REPLACE OLD DEFINITION? Y/(N):y
ANSWER SET L3 HAS BEEN SAVED AS 'RAWMALON8S/A'
```

```
=> e diethylmalonate/a
'A' IS NOT A VALID EXPAND FIELD CODE FOR FILE 'REGISTRY'
The indicated field code is not available for EXPAND in this
file. To see a list of valid EXPAND field codes, enter HELP
SFIELDS at an arrow prompt (=>).
```

```
=> e diethylmalonate/cn
E1      1      DIETHYLMALEIC ANHYDRIDE/CN
E2      1      DIETHYLMALEIMIDE/CN
E3      0 -->  DIETHYLMALONATE/CN
E4      1      DIETHYLMALONATE-FORMALDEHYDE-2,2'-(ISOPROPYLIDENE)BIS(P-PHENY
LENEOXY))DIETHANOL COPOLYMER/CN
E5      1      DIETHYLMALONIC ACID/CN
E6      1      DIETHYLMALONIC ACID DIAMIDE/CN
E7      1      DIETHYLMALONIC ACID DICHLORIDE/CN
E8      1      DIETHYLMALONIC ACID DIETHYL ESTER/CN
E9      1      DIETHYLMALONODINITRILE/CN
E10     1      DIETHYLMALONOHYDRAZIDE/CN
E11     1      DIETHYLMALONONITRILE/CN
E12     1      DIETHYLMALONURIC ACID/CN
```

```
=> e diethyl malonate/cn
E1      1      DIETHYL MALEATE-VINYL CHLORIDE COPOLYMER/CN
E2      1      DIETHYL MALEATE-VINYL CHLORIDE POLYMER/CN
E3      1 -->  DIETHYL MALONATE/CN
E4      1      DIETHYL MALONATE ANION/CN
E5      1      DIETHYL MALONATE BARIUM SALT/CN
E6      1      DIETHYL MALONATE CALCIUM SALT/CN
E7      1      DIETHYL MALONATE ETHOXYMAGNESIUM SALT/CN
E8      1      DIETHYL MALONATE LITHIUM SALT/CN
E9      1      DIETHYL MALONATE SODIUM SALT/CN
E10     1      DIETHYL MALONATE-1,2,3-13C/CN
E11     1      DIETHYL MALONATE-1,2-13C/CN
E12     1      DIETHYL MALONATE-1,3-14C/CN
```

```
=> e3
L4      1      "DIETHYL MALONATE"/CN
```

```
=> e dimethyl malonate/cn
E1      1      DIMETHYL MALEIC ANHYDRIDE-VINYLPYRROLIDINE COPOLYMER/CN
E2      1      DIMETHYL MALEIMIDOMETHYLPHOSPHONATE/CN
E3      1 -->  DIMETHYL MALONATE/CN
E4      1      DIMETHYL MALONATE ANION/CN
E5      1      DIMETHYL MALONATE ION(1-)/CN
E6      1      DIMETHYL MALONATE LITHIUM SALT/CN
E7      1      DIMETHYL MALONATE POTASSIUM SALT/CN
E8      1      DIMETHYL MALONATE SODIUM SALT/CN
E9      1      DIMETHYL MALONATE(1-)/CN
E10     1      DIMETHYL MALONATE-1,1'-(1,2-ETHANEDIYL)BIS(2,2,6,6-TETRAMETH
YL-4-PIPERIDINOL) COPOLYMER/CN
E11     1      DIMETHYL MALONATE-1,1'-(1,2-ETHANEDIYL)BIS(2,2,6,6-TETRAMETH
```

YL-4-PIPERIDINOL) COPOLYMER, SRU/CN
E12 1 DIMETHYL MALONATE-1,3-PROPANEDIOL COPOLYMER/CN

=> e3
L5 1 "DIMETHYL MALONATE"/CN

=> file caplus	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	206.95	208.05

FILE 'CAPLUS' ENTERED AT 08:54:57 ON 16 MAR 2010
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Mar 2010 VOL 152 ISS 12
FILE LAST UPDATED: 15 Mar 2010 (20100315/ED)
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Dec 2009
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Dec 2009

Caplus now includes complete International Patent Classification (IPC) reclassification data for the first quarter of 2010.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> 13
L6 376 L3

=> 13/prep
376 L3
4951107 PREP/RL
L7 305 L3/PREP
(L3 (L) PREP/RL)

=> 14
L8 10515 L4

=> 15
L9 5175 L5

=> 18 or 19
L10 14701 L8 OR L9

=> 16 and 110
L11 129 L6 AND L10

=> optical?
L12 1242311 OPTICAL?

=> l11 and l12
L13 6 L11 AND L12

=> d l13 1-6 ti fbib abs

L13 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN
TI Process for preparation of chiral β -bis-substituted aldehydes
AN 2008:843167 CAPLUS
DN 149:200322
TI Process for preparation of chiral β -bis-substituted aldehydes
IN Ma, Dawei; Ma, Anqi
PA Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences,
Peop. Rep. China
SO Faming Zhuanli Shenqing Gongkai Shuomingshu, 14pp.
CODEN: CNXXEV
DT Patent
LA Chinese
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CN 101215236	A	20080709	CN 2007-10173705 CN 2007-10173705	20071228 20071228

OS CASREACT 149:200322; MARPAT 149:200322
AB This invention provides an enantioselective process for the preparation of chiral β -bis-substituted aldehydes with general formula of $R_1CH[CH(CO_2R_2)2]CH_2CHO$ [wherein $R_1 = 2$ -furyl, p -F-C₆H₄, o -Br-C₆H₄, p -NO₂-C₆H₄, alkyl, or alkenyl; $R_2 = Me$ or Bn] comprising Michael addition of α,β -unsatd. aldehydes with malonates in the presence of (R)- or (S)-2-[(diphenyl(trimethylsilyloxy)methyl]-pyrrolidine. For example, (E)-4-fluorocinnamaldehyde was reacted di-Me malonate in water in the presence of (S)-catalyst to give di-Me 2-[(1R)-1-(4-fluorophenyl)-3-oxopropyl]-malonate with 96% e.e. (67%). The process has advantages of high yield and optical purity. The compds. can be used as important intermediate for synthesizing new compds. and medicines.

L13 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN
TI Method for the production of optically active 3-alkylcarboxylic acids and their intermediates
AN 2005:128812 CAPLUS
DN 144:36149
TI Method for the production of optically active 3-alkylcarboxylic acids and their intermediates
IN Sorger, Klas; Stohrer, Juergen
PA Consortium Fuer Elektrochemische Industrie GmbH, Germany
SO PCT Int. Appl., 52 pp.
CODEN: P1XXD2
DT Patent
LA German
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005115955	A1	20051208	WO 2005-EP52163	20050512
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,				

SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU,
 ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT,
 RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
 MR, NE, SN, TD, TG

DE 102004025901	A1	20051222	DE 2004-102004025901A	20040527
EP 1748975	A1	20070207	DE 2004-102004025901	20040527
EP 1748975	B1	20080213	EP 2005-748062	20050512
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
AT 386010	T	20080315	DE 2004-102004025901A	20040527
			WO 2005-EP52163	W 20050512
			AT 2005-748062	20050512
ES 2299036	T3	20080516	DE 2004-102004025901A	20040527
			ES 2005-748062	20050512
			DE 2004-102004025901A	20040527
US 20070225519	A1	20070927	US 2007-569452	20070329
US 7534908	B2	20090519		
			DE 2004-102004025901A	20040527
			WO 2005-EP52163	W 20050512

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS CASREACT 144:36149; MARPAT 144:36149

AB An enantioselective method for producing optically active 3-alkylcarboxylic acids [e.g., (R)-3-methylheptanoic acid] comprises: (A) an optically active secondary alc. [e.g., (S)-2-hexanol] is transformed into an optically active, activated compound by introducing a terminal group; (B) the activated compound is reacted with a malonic acid derivative so as to obtain an optically active, alkylated malonic acid compound, the reaction taking place exclusively in the presence of one or several solvents selected from ethers or carboxylate esters and one or several aprotic polar solvents or alcs. being optionally added as a cosolvent at a maximum proportion of 30% of the total added solvent volume, provided that the added cosolvent is not hexamethyl phosphoric acid triamide; (C) the malonic acid compound [e.g., [(R)-1-methylpentyl]malonic acid di-Et ester] is saponified if necessary to obtain the corresponding acid; and (D) the corresponding acid [e.g., [(R)-1-methylpentyl]malonic acid] is finally decarboxylated.

RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN

TI Preparation of optically-active
 N-benzyl-5,6-dehydro-3-methylpiperidones as drug intermediates
 AN 1999:481301 CAPLUS
 DN 131:129904
 TI Preparation of optically-active
 N-benzyl-5,6-dehydro-3-methylpiperidones as drug intermediates
 IN Kobayashi, Kaoru; Kusuda, Shinya
 PA Ono Pharmaceutical Co., Japan
 SO Jpn. Kokai Tokkyo Koho, 10 pp.
 CODEN: JKXXAF

DT Patent
 LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 11209345	A	19990803	JP 1998-11035	19980123
				JP 1998-11035	19980123

OS CASREACT 131:129904; MARPAT 131:129904

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The title compds. I [R = CH₂C₆H₄OMe-4, CH₂C₆H₃(OMe)₂-3,4, CH₂C₆H₄Ph-4, CPh₃, CH₂C₆H₄Cl-4, CH₂Ph] or their isomers II, useful as intermediates for condensed piperidine compds. as NO synthase inhibitors, are prepared by dehydration of hydroxypiperidones III (R = same as above) or their isomers IV, resp. III or IV may be prepared by amidation of Me (R)-5-hydroxy-3-methylpentanoate or (R)-4-methyltetrahydro-2H-pyran-2-one, oxidative cyclization of the resulting (R)- or (S)-HOCH₂CH₂CHMeCH₂CONHR (R = same as above), resp. Me (R)-5-hydroxy-3-methylpentanoate may be prepared by reducing Me (R)-3-methylglutarate. (R)-4-methyltetrahydro-2H-pyran-2-one may be prepared by conversion of Me (R)-3-methylglutarate to alkali metal salts, reduction of the salts, and lactonization. Me (R)-3-methylglutarate, prepared by treatment of di-Me 3-methylglutarate (preparation given) with porcine liver esterase, was treated with BH₃.Me₂S in THF at ≤10° for 15 min and then at 25-30° for 1 h to give 97% Me (R)-5-hydroxy-3-methylpentanoate. Me (R)-5-hydroxy-3-methylpentanoate was treated with 4-MeOC₆H₄CH₂NH₂ in toluene under reflux for 1.5 h, and after removing a part of toluene containing low-boiling matters, further refluxed for 2.5 h to give 100% (R)-N-(4-methoxybenzyl)-5-hydroxy-3-methylvaleramide. A DMSO solution of the amide was treated with Et₃N and SO₃-pyridine complex at 15-20° for 30 min to give III (R = CH₂C₆H₄OMe-4), which was treated with p-MeC₆H₄SO₃H in toluene under azeotropic removal of H₂O to give 92.5% I (R = CH₂C₆H₄OMe-4).

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L13 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN

TI Asymmetric synthesis of steroids. XII. Synthesis of the optically active 7α(β),18-dimethyl-19-nortestosterone

AN 1986:186692 CAPLUS

DN 104:186692

OREF 104:29569a,29572a

TI Asymmetric synthesis of steroids. XII. Synthesis of the optically active 7α(β),18-dimethyl-19-nortestosterone

AU Zhuang, Zhiping; Zhou, Weishan

CS Shanghai Inst. Org. Chem., Acad. Sin., Shanghai, Peop. Rep. China

SO Huaxue Xuebao (1985), 43(8), 798-9


CODEN: HHHPA4; ISSN: 0567-7351

DT Journal


LA Chinese

OS CASREACT 104:186692

GI



I



II

AB The title compds. (I) were prepared from oxohexenylcyclopentanone II via

Grignard reaction with m-MeOC6H4CH2Cl, cyclization, redns., and then hydrolysis. Thin-layer chromatog. separation of the mixture gave optically active 7 α - and 7 β -I.

- L13 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN
TI Stereoselective reactions. II. Asymmetric synthesis of β -substituted aldehydes by Michael reaction using chiral α,β -unsaturated aldimines
AN 1980:75775 CAPLUS
DN 92:75775
OREF 92:12475a,12478a
TI Stereoselective reactions. II. Asymmetric synthesis of β -substituted aldehydes by Michael reaction using chiral α,β -unsaturated aldimines
AU Hashimoto, Shunichi; Komeshima, Nobuyasu; Yamada, Shunichi; Koga, Kenji
CS Fac. Pharm. Sci., Univ. Tokyo, Tokyo, 113, Japan
SO Chemical & Pharmaceutical Bulletin (1979), 27(10), 2437-41
CODEN: CPBTAL; ISSN: 0009-2363
DT Journal
LA English
AB The Michael reaction of di-Et malonate with chiral MeCH:CHCH:NCHRCO2CMe3 (R = Me2CH, Me2CHCH2, Me3C), prepared from crotonaldehyde and optically active H2NCHRCO2CMe3 gave the corresponding CHCCH2CHMeCH(CO2Et)2 in reasonably high optical yields after hydrolysis. A proposed stereochem. mechanism of the reaction is presented.
- L13 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN
TI Hydroformylation of some optically active olefins
AN 1975:85633 CAPLUS
DN 82:85633
OREF 82:13691a,13694a
TI Hydroformylation of some optically active olefins
AU Piacenti, F.; Bianchi, M.; Frediani, P.
CS Univ. Firenze, Florence, Italy
SO Advances in Chemistry Series (1974), 132(Homogeneous Catal.-2, Symp., 1973), 283-94
CODEN: ADCSAJ; ISSN: 0065-2393
DT Journal
LA English
AB The hydroformylation of several olefins in the presence of Co2(CO)8 under high CO pressure was examined (S)-5-methylheptanal (75%) and (S)-3-ethylhexanal (4.8%) were products from (+)(S)-4-methyl-2-hexene with optical yields of 94 and 72%, resp. The main products from (+)(S)-2,2,5-trimethyl-3-heptene were (S)-3-ethyl-6,6-dimethylheptanal (56.6%) and (R)-4,7,7-trimethyloctanal (41.2%) obtained with optical yields of 74 and 62%, resp. (R)(S)-3-ethyl-6,6-dimethylheptanal (3.5%) and (R)(S)-4,7,7-trimethyloctanal (93.5%) were formed from (R)(S)-3,6,6-trimethyl-1-heptene. (+)(S)-1-Phenyl-3-methyl-1-pentene, under oxo conditions, was almost completely hydrogenated to (+)(S)-1-phenyl-3-methylpentane with 100% optical yield. 3-(Methyl-d3)-1-butene-4,4,4-d3 gave 4-(methyl-d3)pentanal-5,5,5-d3 (92%), 2-methyl-3-(methyl-d3)butanal-4,4,4-d3 (3.7%), 3-(methyl-d3)pentanal-2,2,d2,3-d (4.3%) with practically 100% retention of D. The reaction mechanism was discussed.
- OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

=>

=> logoff hold

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

FULL ESTIMATED COST	ENTRY 46.72	SESSION 254.77
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-5.10	-5.10

SESSION WILL BE HELD FOR 120 MINUTES
STN INTERNATIONAL SESSION SUSPENDED AT 09:23:22 ON 16 MAR 2010

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1623PAZ

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1		Web Page for STN Seminar Schedule - N. America
NEWS 2	AUG 10	Time limit for inactive STN sessions doubles to 40 minutes
NEWS 3	AUG 18	COMPENDEX indexing changed for the Corporate Source (CS) field
NEWS 4	AUG 24	ENCOMPLIT/ENCOMPLIT2 reloaded and enhanced
NEWS 5	AUG 24	CA/Caplus enhanced with legal status information for U.S. patents
NEWS 6	SEP 09	50 Millionth Unique Chemical Substance Recorded in CAS REGISTRY
NEWS 7	SEP 11	WPIDS, WPINDEX, and WPIX now include Japanese FTERM thesaurus
NEWS 8	OCT 21	Derwent World Patents Index Coverage of Indian and Taiwanese Content Expanded
NEWS 9	OCT 21	Derwent World Patents Index enhanced with human translated claims for Chinese Applications and Utility Models
NEWS 10	NOV 23	Addition of SCAN format to selected STN databases
NEWS 11	NOV 23	Annual Reload of IFI Databases
NEWS 12	DEC 01	FRFULL Content and Search Enhancements
NEWS 13	DEC 01	DGENE, USGENE, and PCTGEN: new percent identity feature for sorting BLAST answer sets
NEWS 14	DEC 02	Derwent World Patent Index: Japanese FI-TERM thesaurus added
NEWS 15	DEC 02	PCTGEN enhanced with patent family and legal status display data from INPADOCDB
NEWS 16	DEC 02	USGENE: Enhanced coverage of bibliographic and sequence information
NEWS 17	DEC 21	New Indicator Identifies Multiple Basic Patent Records Containing Equivalent Chemical Indexing in CA/Caplus
NEWS 18	JAN 12	Match STN Content and Features to Your Information Needs, Quickly and Conveniently
NEWS 19	JAN 25	Annual Reload of MEDLINE database
NEWS 20	FEB 16	STN Express Maintenance Release, Version 8.4.2, Is Now Available for Download
NEWS 21	FEB 16	Derwent World Patents Index (DWPI) Revises Indexing

of Author Abstracts
 NEWS 22 FEB 16 New FASTA Display Formats Added to USGENE and PCTGEN
 NEWS 23 FEB 16 INPADOCDB and INPAFAMDB Enriched with New Content
 and Features
 NEWS 24 FEB 16 INSPEC Adding Its Own IPC codes and Author's E-mail
 Addresses
 NEWS EXPRESS FEBRUARY 15 10 CURRENT WINDOWS VERSION IS V8.4.2,
 AND CURRENT DISCOVER FILE IS DATED 15 JANUARY 2010.
 NEWS HOURS STN Operating Hours Plus Help Desk Availability
 NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that
 specific topic.

All use of STN is subject to the provisions of the STN customer
 agreement. This agreement limits use to scientific research. Use
 for software development or design, implementation of commercial
 gateways, or use of CAS and STN data in the building of commercial
 products is prohibited and may result in loss of user privileges
 and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 11:49:35 ON 16 MAR 2010

=> off hold

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.22

0.22

SESSION WILL BE HELD FOR 120 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 11:49:41 ON 16 MAR 2010